

# ALGORITHM FOR USE

Patient experiences nausea



Patient peels back blue foil top, leaving white perforated barrier intact



Patient holds just below nose, then takes deep breaths as often and as long as necessary



QueaseEASE is labeled with patient sticker



Patient is allowed to keep QueaseEASE in accessible area, in hand, in a pocket, etc



Patient is instructed to use as often as needed throughout postoperative phase:



Including PACU, car ride home or any time nausea is encountered.



If collecting data for efficacy study, fill in collection tool

# QueaseEASE DATA COLLECTION TOOL

Patient Age \_\_\_\_\_ DOB \_\_\_\_/\_\_\_\_/\_\_\_\_ Gender M F

Procedure \_\_\_\_\_

History of PONV Yes No Motion Sickness Yes No

Smoker Yes No Antiemetics given intraop? Yes No

Was QueaseEASE used as a first line rescue remedy? Yes No

Additional antiemetics given? Yes No

Which statement below represents the patient experience with QueaseEASE the most?

- It took away the nausea completely
- It helped somewhat, but still felt slightly nauseated
- It gave them minor relief, but they still felt nauseated
- It had NO effect on nausea

Did patient feel QueaseEASE was beneficial? Yes No

Was QueaseEASE sent with patient post operatively? Yes No

Patient Stay Time \_\_\_\_\_am/pm

Additional Comments \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_